


# Angomonas deanei

Natascha A. van Geelen-Kuenzel <sup>1</sup>, Vyacheslav Yurchenko <sup>2</sup>, Julius Lukeš <sup>3</sup>, and Eva C.M. Nowack <sup>1,\*</sup>

<sup>1</sup>Institute of Microbial Cell Biology, Heinrich Heine University, Düsseldorf, Germany

<sup>2</sup>Life Science Research Centre, Faculty of Science, University of Ostrava, Ostrava, Czechia

<sup>3</sup>Institute of Parasitology, Biology Centre, Czech Academy of Sciences, and Faculty of Sciences, University of South Bohemia, České Budějovice, Czechia



**KEY FACTS:**

*A. deanei* harbors a vertically transmitted cytoplasmic  $\beta$ -proteobacterial endosymbiont, *Ca. Kinetoplastibacterium crithidii* (Alcaligenaceae), which is found exclusively in the trypanosomatid subfamily Strigomonadinae and is most closely related to the genus *Taylorella* containing pathogens of horses and donkeys.

The 0.8 Mbp genome of *Ca. K. crithidii* is streamlined to support host metabolism. The endosymbiont associates with host glycosomes – metabolically specialized peroxisomes.


The endosymbiont's division machinery has been largely replaced by a host-derived dynamin-based division system.

The high-quality genome assembly of *A. deanei* contains 10 365 protein-coding genes.

*A. deanei* became a versatile, genetically tractable model system for studying the molecular mechanisms underpinning host/endosymbiont interaction.

The species has been originally described as *Crithidia deanei* Carvalho, 1973. Another formally described species, *Herpetomonas roitmani* Fiorini, 1989, is indistinguishable from *A. deanei* by molecular analyses.

*Angomonas deanei* is a cosmopolitan species of monoxenous trypanosomatids parasitizing the gut of numerous insects. It harbors a tightly integrated bacterial endosymbiont that complements the metabolism of its osmotrophic host, supplying heme, several amino acids, nucleotides, and vitamins. The endosymbiont is closely associated with host glycosomes that adapted in protein composition, likely to provide the endosymbiont with metabolic intermediates. Remarkably, the single endosymbiont grows and divides in synchrony with its host. This synchronization of cell cycles appears to be enabled by the loss of most of the bacterial division machinery and evolution of a set of endosymbiont-targeted host proteins (ETPs). A dynamin-based machinery composed of ETPs became essential for endosymbiont division and may reflect an early stage of a trajectory converting a bacterium into an organelle. Efficient genetic manipulation makes *A. deanei* a versatile model for studying host/symbiont interaction.



**DISEASE FACTS:**

*A. deanei* is a parasite or commensal of numerous true bug and fly species.

Its life cycle does not include host switching.

**TAXONOMY AND CLASSIFICATION:**

**PHYLUM:** Euglenozoa  
**CLASS:** Kinetoplastea  
**ORDER:** Trypanosomatida  
**FAMILY:** Trypanosomatidae  
**SUBFAMILY:** Strigomonadinae  
**GENUS:** *Angomonas*  
**SPECIES:** *A. deanei* Carvalho, 1973 emend. Teixeira et Camargo, 2011

\*Correspondence:

[e.nowack@hhu.de](mailto:e.nowack@hhu.de) (E.C.M. Nowack).

<https://doi.org/10.1016/j.pt.2025.12.008> 309



## Acknowledgments

This work was supported by the German Research Foundation (SFB1535, project ID 458090666 to E.C.M.N.), Czech Science Foundation (25-15298S to V.Y. and J.L.), and by the EU via the Operational Programme Just Transition and the Ministry of the Environment (CZ.10.03.01/00/22\_003/0000003 LERCO to V.Y.).

## Declaration of interests

The authors declare no competing interests.

## Resources

[www.ncbi.nlm.nih.gov/datasets/genome/GCA\\_903995115.1/](http://www.ncbi.nlm.nih.gov/datasets/genome/GCA_903995115.1/)  
[www.ncbi.nlm.nih.gov/datasets/genome/GCF\\_000340825.1/](http://www.ncbi.nlm.nih.gov/datasets/genome/GCF_000340825.1/)  
<https://tritypdb.org/>

## Literature

1. Alves, J.M.P. *et al.* (2013) Endosymbiosis in trypanosomatids: the genomic cooperation between bacterium and host in the synthesis of essential amino acids is heavily influenced by multiple horizontal gene transfers. *BMC Evol. Biol.* 13, 190
2. Carvalho, A.L.M. (1973) Estudos sobre a posição sistemática, a biologia e a transmissão de tripanosomatídeos encontrados em *Zelus leucogrammus* (Perty, 1834) (Hemiptera, Reduviidae). *Rev. Pathol. Trop.* 2, 223–274
3. Davey, J.W. *et al.* (2021) Chromosomal assembly of the nuclear genome of the endosymbiont-bearing trypanosomatid *Angomonas deanei*. *G3 Genes Genomes Genet.* 11, jkaa018
4. Maurya, A.K. *et al.* (2025) A nucleus-encoded dynamin-like protein controls endosymbiont division in the trypanosomatid *Angomonas deanei*. *Sci. Adv.* 11, eadp8518
5. Morales, J. *et al.* (2016) Development of a toolbox to dissect host-endosymbiont interactions and protein trafficking in the trypanosomatid *Angomonas deanei*. *BMC Evol. Biol.* 16, 247
6. Morales, J. *et al.* (2023) Host-symbiont interactions in *Angomonas deanei* include the evolution of a host-derived dynamin ring around the endosymbiont division site. *Curr. Biol.* 33, 28–40
7. Motta, M.C. *et al.* (2010) The bacterium endosymbiont of *Crithidia deanei* undergoes coordinated division with the host cell nucleus. *PLoS One* 5, e12415
8. Skalický, T. *et al.* (2021) Endosymbiont capture, a repeated process of endosymbiont transfer with replacement in trypanosomatids *Angomonas* spp. *Pathogens* 10, 702
9. Teixeira, M.M.G. *et al.* (2011) Phylogenetic validation of the genera *Angomonas* and *Strigomonas* of trypanosomatids harboring bacterial endosymbionts with the description of new species of trypanosomatids and of proteobacterial symbionts. *Protist* 162, 503–524
10. Yurchenko, V. *et al.* (2009) Selective recovery of the cultivation-prone components from mixed trypanosomatid infections: a case of several novel species isolated from Neotropical Heteroptera. *Int. J. Syst. Evol. Microbiol.* 59, 893–909