

ON THE UTILITY OF ROBINSON–AMITSUR ULTRAFILTERS

PASHA ZUSMANOVICH

ABSTRACT. An embedding theorem for algebraic systems is presented, basing on a certain old ultrafilter construction. As an application, we outline alternative proofs of some results from the theory of PI algebras, and establish some properties of Tarski’s monsters.

INTRODUCTION

In 1960s, A. Robinson and S. Amitsur established a number of embedding results in Ring Theory which proved to be useful in various structural questions. A typical example: if a prime ring R embeds in a direct product of associative division rings, then R embeds in an associative division ring (see [Am1, Proof of Theorem 15] and [Am2, Theorem 3] for original papers, and [E, §6] for a nice overview).

The proof of these results follows the same scheme: basing on the initial data – a ring embedded in a direct product of rings – a certain sort of ultrafilter, which we call a *Robinson–Amitsur ultrafilter*, is constructed. Using this ultrafilter, one passes from the direct product of rings to their ultraproduct, and appeal to the Łoś theorem about elementary equivalence of an algebraic system and its ultraproduct completes the proof.

In this paper we extend this argument to a class of general algebraic systems (Theorem 1.1), and observe similarity with the celebrated Jónsson lemma from the universal algebra. Coupled with the classical Birkhoff theorem about varieties of algebraic systems, this gives a simple yet elegant criterion for an algebra or group not to satisfy a nontrivial identity (Corollaries 2.5 and 3.2). The corresponding group result is not entirely new (see comments after Corollary 3.2), but, we believe, its proof is, and the links between apriori unrelated concepts, ideas and results is the main novelty of this paper.

As an application, we outline alternative, “by abstract nonsense”, proofs of some particular cases of well-known results from the theory of PI algebras (§4), of results about algebras having the same identities (§5), and establish that Tarski’s monsters without identities have infinite (relative) girth (§6). Finally, in §7 we formulate and prove a “dual” version of Theorem 1.1, with embeddings replaced by surjective homomorphisms, and discuss some its consequences and related questions.

The narrative is occasionally interspersed with questions and speculations.

1. ALGEBRAIC SYSTEMS

In what follows, by an *algebra* or *ring*, we mean an arbitrary, not necessarily associative, or Lie, or satisfying any other distinguished identities, algebra or ring, unless it is stated otherwise. Algebras are considered over fields. *Ideal* of an algebra means a two-sided ideal. By an (*algebraic*) *system* we mean an algebra in the universal algebraic sense, i.e. a set with a number of operations on it of, generally, various arity.

We deal with algebraic systems whose congruences behave “good enough”, like ideals in rings, or normal subgroups in groups (or, more generally, ideals in the so-called Ω -groups introduced by P.J. Higgins and studied by Kurosh and his school (see [Ku, Chapter 3, §2])). Namely, suppose the signature Ω of a class of algebraic systems has a 0-ary operation e (i.e., a distinguished element). A term $t(x_1, \dots, x_n, y_1, \dots, y_m)$ composed of the operations in Ω is called an *ideal term* in y_1, \dots, y_m if $m > 0$ and

$$t(a_1, \dots, a_n, e, \dots, e) = e$$

Date: last (cosmetical) revision September 16, 2025.

J. Algebra **388** (2013), 268–286.

for any $a_1, \dots, a_n \in A$ for any system A in the class. A subset I of an algebraic system A is called an *ideal* if for any ideal term $t(x_1, \dots, x_n, y_1, \dots, y_m)$ in y_1, \dots, y_m , the element

$$(1) \quad t(a_1, \dots, a_n, b_1, \dots, b_m)$$

belongs to I for any $a_1, \dots, a_n \in A$ and $b_1, \dots, b_m \in I$.

For every congruence θ on A , its equivalence class $\{a \in A \mid a \equiv_\theta e\}$ is an ideal of A . Generally, this relation between ideal and congruences is not one-to-one, but if it is, the corresponding class of algebraic systems is called *ideal-determined*. In particular, for systems from an ideal-determined class we can speak about quotients by ideals instead of quotients by congruences.

This notion, along with its numerous particular cases and variations, was studied by Agliano, Chajda, Fichtner, Grätzer, Gumm, Słomiński, Ursini and others (see, for example, [CEL, Chapter 10]).

Any algebraic system A from an ideal-determined class has at least two ideals – a *trivial ideal* $\{e\}$ and the whole A . The intersection of any two ideals of A is an ideal. A is called *finitely subdirectly irreducible* if intersection of any two its nontrivial ideals is nontrivial. An *ideal generated by the subset* X of A is the minimal ideal of A containing X , and it coincides with the set of all elements of the form (1), where $t(x_1, \dots, x_n, y_1, \dots, y_m)$ is an ideal term in y_1, \dots, y_m , and $a_1, \dots, a_n \in A$, $b_1, \dots, b_m \in X$.

Recall the construction of the ultraproduct – not in the most general form, but in the form suitable for our purposes. Let $\{A_i\}_{i \in \mathcal{I}}$ be a set of algebraic systems from an ideal-determined class, and \mathcal{F} a filter on the indexing set \mathcal{I} . Then

$$\mathcal{I}\left(\prod_{i \in \mathcal{I}} A_i, \mathcal{F}\right) = \{f \in \prod_{i \in \mathcal{I}} A_i \mid \{i \in \mathcal{I} \mid f(i) = e\} \in \mathcal{F}\}$$

is an ideal of the direct product $\prod_{i \in \mathcal{I}} A_i$, and the quotient by this ideal is called a *filtered product* of the set $\{A_i\}_{i \in \mathcal{I}}$ with respect to the filter \mathcal{F} , and is denoted by $\prod_{\mathcal{F}} A_i$. In the particular case where all A_i 's are isomorphic to the same algebraic system A , their filtered product is called a *filtered power* of A and is denoted by $A^{\mathcal{F}}$. When \mathcal{F} is an ultrafilter, we speak about *ultraproducts* and *ultrapowers*.

An alternative view on ultrafilters and ultraproducts is an analytic one: ultrafilter is a probability measure taking only two values – 0 and 1, and such that every subset of the indexing set is measurable. An ultrapower is the collection of measurable functions from the indexing set considered up to the measure.

We refer to [CEL], [C], [FM], [Ku], or [M] for all necessary basic notions and results related to universal algebra, and to [BS], [C], [E], or [M] again for ultrafilters and ultraproducts.

Theorem 1.1 (Robinson–Amitsur for algebraic systems). *Let $\{B_i\}_{i \in \mathcal{I}}$ be a set of algebraic systems from an ideal-determined class. If a finitely subdirectly irreducible algebraic system A embeds in the direct product $\prod_{i \in \mathcal{I}} B_i$, then there is an ultrafilter \mathcal{U} on the set \mathcal{I} such that A embeds in the ultraproduct $\prod_{\mathcal{U}} B_i$.*

Proof. Define

$$\mathcal{S} = \{\{i \in \mathcal{I} \mid f(i) \neq e\} \mid f \in A, f \neq e\}.$$

Let us verify that intersection of any two elements of \mathcal{S} contains an element of \mathcal{S} . Let $S, T \in \mathcal{S}$, say, $S = \{i \in \mathcal{I} \mid f(i) \neq e\}$ and $T = \{i \in \mathcal{I} \mid g(i) \neq e\}$ for some $f, g \in A$ different from e . Since A is finitely subdirectly irreducible, it contains an element $u \neq e$ belonging to the intersection of ideals generated by $\{f\}$ and $\{g\}$. Let $i \in \mathcal{I}$ such that $f(i) = e$. Since $u = t(h_1, \dots, h_n, f, \dots, f)$ for some ideal term t and $h_1, \dots, h_n \in A$,

$$\begin{aligned} u(i) &= t(h_1, \dots, h_n, f, \dots, f)(i) = t(h_1(i), \dots, h_n(i), f(i), \dots, f(i)) \\ &= t(h_1(i), \dots, h_n(i), e, \dots, e) = e. \end{aligned}$$

Coupling this with a similar assertion for g , we get that

$$S \cap T \supset \{i \in \mathcal{I} \mid u(i) \neq e\} \in \mathcal{S}.$$

Thus \mathcal{S} satisfies the finite intersection property and is contained in some ultrafilter \mathcal{U} on \mathcal{I} . Factoring the embedding of algebraic systems $A \hookrightarrow \prod_{i \in \mathcal{I}} B_i$ by the ideal $\mathcal{I}(\prod_{i \in \mathcal{I}} B_i, \mathcal{U})$, we get an embedding of

algebraic systems

$$A / \left(A \cap \mathcal{I} \left(\prod_{i \in \mathcal{J}} B_i, \mathcal{U} \right) \right) \hookrightarrow \left(\prod_{i \in \mathcal{J}} B_i \right) / \mathcal{I} \left(\prod_{i \in \mathcal{J}} B_i, \mathcal{U} \right) = \prod_{\mathcal{U}} B_i.$$

Let $f \in A \cap \mathcal{I} \left(\prod_{i \in \mathcal{J}} B_i, \mathcal{U} \right)$. Then $\{i \in \mathcal{J} \mid f(i) = e\} \in \mathcal{U}$, and, since \mathcal{U} is an ultrafilter, $\{i \in \mathcal{J} \mid f(i) \neq e\} \notin \mathcal{U}$, and hence $\{i \in \mathcal{J} \mid f(i) \neq e\} \notin \mathcal{S}$. From $f \in A$ and the definition of \mathcal{S} it follows that $f = e$. This shows that $A \cap \mathcal{I} \left(\prod_{i \in \mathcal{J}} B_i, \mathcal{U} \right) = \{e\}$. \square

The ultraproduct construction used in this proof mimics the old one, used by A. Robinson and S. Amitsur in Ring Theory, mentioned in the introduction.

The finite subdirect irreducibility of an algebraic system A is equivalent to the following condition: if A embeds in the finite direct product of algebraic systems $\prod_{i=1}^n B_i$, then A embeds in one of B_i 's. An infinite analog of this condition is *subdirect irreducibility*, that is, the condition that intersection of any (possibly infinite) set of nontrivial ideals of A is nontrivial (or, equivalently, A possesses a minimal nontrivial ideal which is called *monolith*). Similarly, the latter condition is equivalent to the following: if A embeds in the (possibly infinite) direct product $\prod_{i \in \mathcal{J}} B_i$, then A embeds in one of B_i 's. Thus, Theorem 1.1 can be considered as, perhaps, somewhat surprising statement that for ideal-determined classes, finite subdirect irreducibility implies a sort of a weaker form of subdirect irreducibility.

An application of Theorem 1.1 to varieties and quasivarieties of algebraic systems follows.

If A is an algebraic system, $\text{Var}(A)$ and $\text{Qvar}(A)$ denote, respectively, a variety and a quasivariety generated by A . Any quasivariety (and, in particular, any variety) possesses free algebraic systems (this is formulated explicitly, for example, in [C, Chapter VI, Proposition 4.5] and is implicit in [M, Chapter V]).

Corollary 1.2. *Let A be an algebraic system from an ideal-determined class. A finitely subdirectly irreducible free system in $\text{Var}(A)$ or $\text{Qvar}(A)$ embeds in an ultrapower of A .*

Proof. Let \mathcal{F} be a free system in $\text{Var}(A)$ which is finitely subdirectly irreducible. According to the Birkhoff theorem, $\mathcal{F} = B/I$ for an ideal I of an algebra B , and B is a subalgebra of a direct power of A . Because of the universal property of \mathcal{F} , the short exact sequence $\{e\} \rightarrow I \rightarrow B \rightarrow \mathcal{F} \rightarrow \{e\}$ splits, i.e. \mathcal{F} embeds in B , and hence in a direct power of A . Then apply Theorem 1.1.

Similarly, according to the Birkhoff-like characterization of quasivarieties due to Malcev ([M, Chapter V, §11, Theorem 4]), if \mathcal{F} is a free system in $\text{Qvar}(A)$, then it is a subalgebra of a filtered power $A^{\mathcal{F}} = A^{\mathcal{J}} / \mathcal{I}(A^{\mathcal{J}}, \mathcal{F})$ of A . Taking preimage of \mathcal{F} with respect to the homomorphism $A^{\mathcal{J}} \rightarrow A^{\mathcal{J}} / \mathcal{I}(A^{\mathcal{J}}, \mathcal{F})$, we get that \mathcal{F} is a quotient of a subalgebra in $A^{\mathcal{J}}$, and the rest of reasoning is the same as above. \square

Compare Theorem 1.1 and Corollary 1.2 with the celebrated Jónsson lemma in the generalized form due to combined efforts of Freese, Hagemann, Herrmann, Hrushovski and McKenzie (see [FM, Theorem 10.1]): if A is a subdirectly irreducible algebraic system from a modular variety (i.e., the congruence lattice of any system from the variety is modular) generated by a set $\{B_i\}_{i \in \mathcal{J}}$ of algebraic systems, then the quotient of A by the centralizer of its monolith embeds in a homomorphic image of a subsystem of an ultraproduct $\prod_{\mathcal{U}} B_i$. Note that the congruence (=ideal) lattice of any algebra from an ideal-determined class is modular (see, for example, [CEL, Remark 10.1.16]).

Corollary 1.3 (Criterion for absence of non-trivial identities for algebraic systems). *Let \mathfrak{V} be a variety of algebraic systems from an ideal-determined class, and suppose that all free systems of \mathfrak{V} are finitely subdirectly irreducible. Then for an algebraic system $A \in \mathfrak{V}$, the following is equivalent:*

- (i) *any identity of A is an identity of \mathfrak{V} (i.e., A does not satisfy nontrivial identities within \mathfrak{V});*
- (ii) *any free system of \mathfrak{V} embeds in an ultrapower of A ;*
- (iii) *any free system of \mathfrak{V} embeds in a system elementarily equivalent to A .*

Proof. (i) \Rightarrow (ii) follows from Corollary 1.2.

(ii) \Rightarrow (iii) follows from the Łoś theorem about elementary equivalence of an algebraic system and its ultrapower.

(iii) \Rightarrow (i) An algebraic system elementarily equivalent to A does not satisfy a nontrivial identity within \mathfrak{V} . Since the latter is the first-order property, A does not satisfy a nontrivial identity either. \square

Remarks.

(1) Of course, the equivalence of conditions (ii) and (iii) also follows from the (powerful) Keisler ultrapower theorem (see, for example, [BS, Chapter 7, Corollary 2.7]).

(2) Recall a well-known fact from model theory: an algebraic system B embeds in an ultrapower of an algebraic system A if and only if $Th_{\forall}(A) \subseteq Th_{\forall}(B)$, where Th_{\forall} denotes the universal theory of a system (see, for example, [BS, Chapter 9, Lemma 3.8]). This allows to rephrase condition (ii) or (iii) as follows: the universal theory of A is contained in the universal theory of any free system of \mathfrak{V} .

(3) As any variety is determined by its free system of countable rank (see, for example, [C, Chapter IV, Proposition 3.8] or [M, Chapter VI, §13, Theorem 3]), in conditions (ii) and (iii) of the corollary, as well as in (2) above, one may replace “any free system” by “the free system of countable rank”.

2. ALGEBRAS

Specializing results of the previous section to the (ideal-determined, obviously) class of rings, we get, for example, a nonassociative analog of one of the classical embedding results in Ring Theory mentioned in the introduction:

Corollary 2.1. *If a finitely subdirectly irreducible ring A embeds in a direct product of division rings, then A embeds in a division ring.*

Proof. By Theorem 1.1, A embeds in an ultraproduct of division rings. As the property to be a division ring is the first-order property, by the Łoś theorem the ultraproduct of division rings is a division ring, whence the conclusion. \square

When trying to specialize to the class of algebras, we should deal with the issue of the base field. For example, Corollary 1.3 is not applicable directly. The problem is this: Corollary 1.3 is obtained by combination of the Birkhoff theorem about varieties of algebras (or a similar statements), and the Łoś theorem about elementary equivalence of an algebra and its ultrapower. One may treat algebras either as two-sorted theories (algebra over a field, field), or distinguish elements of the base field by an unary predicate. Either way, while in Birkhoff-like reasonings the base field remains the same, in Łoś ones it, generally, changes: we pass to an ultraproduct of the base field.

So, for now on, fix the base field K . The embedding claimed in Theorem 1.1 is an embedding of algebras defined over K . Of course, the ultraproduct $\prod_{\mathcal{U}} B_i$ is defined also over the ultrapower field $K^{\mathcal{U}}$, so we have an embedding of $K^{\mathcal{U}}A$ in $\prod_{\mathcal{U}} B_i$ as $K^{\mathcal{U}}$ -algebras. (Here $K^{\mathcal{U}}A$ is understood as a $K^{\mathcal{U}}$ -linear span of A , considered as a K -subalgebra of $\prod_{\mathcal{U}} B_i$). Due to the universal property of the tensor product, there is a surjection of $K^{\mathcal{U}}$ -algebras

$$(2) \quad A \otimes_K K^{\mathcal{U}} \rightarrow K^{\mathcal{U}}A,$$

but this surjection is, generally, not a bijection.

An important observation is that for free algebras A in the major varieties of algebras considered in the literature – the varieties of all algebras, associative algebras, and Lie algebras, the surjection (2) is a bijection.

Lemma 2.2. *Let A be a subalgebra of an algebra B , both defined over a field K . Suppose B is also defined over a field F containing K , and that as a K -algebra, A does not have commutative subspaces (i.e., subspaces all whose elements commute) of dimension > 1 . Then $FA \simeq A \otimes_K F$ as F -algebras.*

Proof. The claimed isomorphism follows from the fact that the linear dependence of elements of A over F implies their linear dependence over K . Indeed, consider the linear dependence

$$(3) \quad f_1a_1 + \cdots + f_na_n = 0,$$

where $f_i \in F$, $a_i \in A$, and let us prove by induction on n that a_1, \dots, a_n are linearly dependent over K . For $n = 1$ this is trivial. Taking the commutator with a_n of both sides of (3), we get

$$f_1[a_1, a_n] + \dots + f_{n-1}[a_{n-1}, a_n] = 0.$$

(we use the usual notation for the commutator: $[a, b] = ab - ba$). By inductive hypothesis, there are $k_1, \dots, k_{n-1} \in K$, not all zero, such that

$$k_1[a_1, a_n] + \dots + k_{n-1}[a_{n-1}, a_n] = [k_1 a_1 + \dots + k_{n-1} a_{n-1}, a_n] = 0.$$

But since A does not have commutative subspaces of dimension > 1 ,

$$k(k_1 a_1 + \dots + k_{n-1} a_{n-1}) + \ell a_n = 0$$

for some $k, \ell \in K$, not both zero. □

Obviously, the hypothesis of this lemma is satisfied when A is a free algebra or a free Lie algebra of rank > 1 (in view of the Shirshov–Witt theorem). A similar, but just a little bit more involved argument (see, for example, proof of Lemma 1 in [MM]) shows that the conclusion of the lemma holds also when A is a free associative algebra of rank > 1 .

Another issue we should deal with when trying to apply results of the previous section to the class of algebras, is when the hypothesis of Corollary 1.2 holds, i.e., when free algebras in a variety are finitely subdirectly irreducible. Again, this hypothesis is satisfied for absolutely free algebras, free associative algebras, and free Lie algebras of rank > 1 (on the other hand, it is not satisfied for free Jordan algebras and free alternative algebras). In fact, those free algebras satisfy a stronger condition – primeness, and, in the context of algebras, we will mainly focus on the latter condition instead of finite subdirect irreducibility.

Let \mathfrak{V} be a variety of algebras, and $\mathcal{F}(X)$ is the free algebra in this variety generated by a set X . By *words* we mean elements of $\mathcal{F}(X)$ for some X . The standard grading on $\mathcal{F}(X)$ is defined by length of words. By *2-nontrivial words* we mean words degrees of all whose homogeneous components in each of the first two indeterminates are non zero. For example,

$$xy, \quad (xy)x - (xy)(xz), \quad (zy)x + 2(tx)y + x(yz)(tx)$$

are 2-nontrivial words in x, y, z, t (in that order), while

$$x, \quad xz, \quad (xy)x + xz, \quad (yz)(yt) + (xy)z$$

are not.

Lemma 2.3. *For an algebra $A \in \mathfrak{V}$, the following is equivalent:*

- (i) *For any two nonzero ideals I, J of A , $IJ \neq 0$.*
- (ii) *For any two nonzero elements $x, y \in A$, there is a 2-nontrivial word $w(\xi_1, \dots, \xi_n)$, $n \geq 2$ and elements $x_1, \dots, x_{n-2} \in A$ such that $w(x, y, x_1, \dots, x_{n-2}) \neq 0$.*

Proof. (i) \Rightarrow (ii). Suppose there are nonzero $x, y \in A$ such that for any 2-nontrivial word $w(\xi_1, \dots, \xi_n)$ and any $x_1, \dots, x_{n-2} \in A$, $w(x, y, x_1, \dots, x_{n-2}) = 0$. Let I and J be ideals of A generated by x and y respectively. Clearly $IJ = 0$, a contradiction.

(ii) \Rightarrow (i). Suppose I, J are two nonzero ideals of A . Taking $x \in I$ and $y \in J$, we have $w(x, y, x_1, \dots, x_{n-2}) \neq 0$ for some 2-nontrivial word w and elements x_i of A . Clearly, $w(x, y, x_1, \dots, x_{n-2}) \in I \cap J$.

Further, there is a 2-nontrivial word u and elements $y_1, \dots, y_{m-2} \in A$ such that

$$u(w(x, y, x_1, \dots, x_{n-2}), w(x, y, x_1, \dots, x_{n-2}), y_1, \dots, y_{m-2}) \neq 0.$$

In particular, for some monomial m occurring in u , we have

$$(4) \quad m(w(x, y, x_1, \dots, x_{n-2}), w(x, y, x_1, \dots, x_{n-2}), y_1, \dots, y_{m-2}) \neq 0.$$

Examining how m is built up from the variables, there must be some point where a sub-monomial containing the first variable is multiplied by a sub-monomial containing the second variable, giving a product pq where p involves one of the first two variables, and q the other.

Due to (4),

$$p(w(x, y, x_1, \dots, x_{n-2}), w(x, y, x_1, \dots, x_{n-2}), y_1, \dots, y_{m-2}) \\ \times q(w(x, y, x_1, \dots, x_{n-2}), w(x, y, x_1, \dots, x_{n-2}), y_1, \dots, y_{m-2}) \neq 0.$$

Each of these factors belongs to $I \cap J$ because $w(x, y, x_1, \dots, x_{n-2})$ does, so their product belongs to $(I \cap J)(I \cap J) \subseteq IJ$, as required. \square

An algebra $A \in \mathfrak{V}$ satisfying the equivalent conditions of Lemma 2.3, is called \mathfrak{V} -prime. When \mathfrak{V} is a variety of all associative algebras, this notion coincides with the classical notion of a prime associative algebra.

Clearly, if \mathfrak{W} is another variety and $\mathfrak{V} \subseteq \mathfrak{W}$, an algebra $A \in \mathfrak{V}$ is \mathfrak{V} -prime if and only if it is \mathfrak{W} -prime, so we can simply speak about *prime algebras* (which are prime in the variety of all algebras).

As for any two ideals I and J , $IJ \subseteq I \cap J$, prime algebras (or rings) are finitely subdirectly irreducible.

Now, the claim about primeness of free algebras and free associative algebras is obvious, as they do not have zero divisors. To establish primeness of a free Lie algebra of rank > 1 , consider two nonzero ideals I, J in it. By the Shirshov–Witt theorem about freeness of subalgebras of a free Lie algebra, neither of I, J can be one-dimensional, and hence we may choose two linearly independent elements $x \in I$ and $y \in J$. Their commutator is nonzero, as otherwise they would form a 2-dimensional abelian subalgebra, which again contradicts the Shirshov–Witt theorem. Hence $[I, J] \neq 0$.

Another situation when relatively free algebras are prime is described by the following lemma.

Lemma 2.4. *Let A be a prime algebra over an infinite field. Then the free algebra of infinite rank in $\text{Var}(A)$ is prime.*

Proof. Let $\mathcal{F}(X)$ be the free algebra in $\text{Var}(A)$ freely generated by an infinite set $X = \{x_1, x_2, \dots\}$. Suppose that there are nonzero elements $u(x_1, \dots, x_n), v(x_1, \dots, x_n) \in \mathcal{F}(X)$ such that

$$(5) \quad w(u, v, u_1, \dots, u_m) = 0$$

for any 2-nontrivial word w and $u_1, \dots, u_m \in \mathcal{F}(X)$. As any relation between free generators of $\mathcal{F}(X)$ is an identity in A , the equalities (5) are identities in A . Taking $u_1 = x_{n+1}, u_2 = x_{n+2}, \dots$, we get that

$$w(u(x_1, \dots, x_n), v(x_1, \dots, x_n), x_{n+1}, \dots, x_{n+m}) = 0$$

for any 2-nontrivial word w and any $x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m} \in A$. As A is prime, either $u(x_1, \dots, x_n) = 0$ or $v(x_1, \dots, x_n) = 0$ for any $x_1, \dots, x_n \in A$.

Fix some basis of A , take $(x_1, \dots, x_n) \in A^n = A \times \dots \times A$ (n times) at which u is nonzero, $(y_1, \dots, y_n) \in A^n$ at which v is nonzero, and consider the line in A^n of all affine linear combinations

$$(6) \quad \lambda(x_1, \dots, x_n) + (1 - \lambda)(y_1, \dots, y_n),$$

where λ is an element of the base field K . On this line, the coordinates of u and v in terms of the chosen basis are each given by a polynomial in λ . So, picking a K -linear function on A such whose value at $u(x_1, \dots, x_n)$ is nonzero and another such whose value at $v(y_1, \dots, y_n)$ is nonzero, we see that for all but finitely many $\lambda \in K$, both u and v will be nonzero on the n -tuple (6), a contradiction.

If A is finite-dimensional, we can use a similar, but more elegant Zariski-topology argument instead: both sets of n -tuples of elements of A on which u , respectively v , does not vanish, form a nonempty Zariski-open subset in A^n , whence they have a nonzero intersection, a contradiction. \square

All these observations lead to a variant of the general Corollary 1.3 for the case of algebras:

Corollary 2.5 (Criterion for absence of non-trivial identities for algebras). *For an algebra A belonging to one of the following varieties of algebras: all algebras, associative algebras, or Lie algebras, the following are equivalent:*

- (i) *A does not satisfy a nontrivial identity;*
- (ii) *any free algebra embeds in an ultrapower of A;*
- (iii) *any free algebra embeds in an algebra elementarily equivalent to A.*

Proof. (i) \Rightarrow (ii) follows from Corollary 1.2. Note that in view of Lemma 2.2 (applied to the case $F = K^{\mathcal{U}}$, where K is the base field, and \mathcal{U} is an appropriate ultrafilter) and remarks after it, the embedding of Corollary 1.2 can be considered as an embedding of $K^{\mathcal{U}}$ -algebras.

(ii) \Rightarrow (iii) as in the proof of Corollary 1.3: by the Łoś theorem, the pairs (B, K) and $(B^{\mathcal{U}}, K^{\mathcal{U}})$ are elementarily equivalent as models of the two-sorted theory (algebra over a field, field).

(iii) \Rightarrow (i) as in the proof of Corollary 1.3. \square

All the remarks after Corollary 1.3 are applicable. In addition, as the free associative, respectively Lie, algebra of countable rank embeds in the free associative, respectively Lie, algebra of rank 2, for these varieties one can replace “any free algebra” in conditions (ii) and (iii) by “the free algebra of rank 2”.

Let us provide some negative examples showing that conclusions of some statements of this section do not always hold.

The conclusion of Lemma 2.2 does not hold for polynomial algebras in > 1 variables (i.e., for free associative commutative algebras). Indeed, let $F = K(x)$ (the field of rational functions in 1 variable) and $A = K[x, y]$ (the algebra of polynomials in 2 variables), considered as a K -subalgebra of $K(x, y)$ (the field of rational functions in 2 variables). Then

$$FA = K(x)[y] \simeq K[y] \otimes_K K(x),$$

but $A \otimes_K F = K[x, y] \otimes_K K(x)$.

In some other situations, the surjection (2) also can be very far from being a bijection. For example, consider the free algebra \mathcal{F} of countable rank in the variety $\text{Var}(A)$ generated by a finite-dimensional prime algebra over an infinite field K . By Lemma 2.4, \mathcal{F} is prime, and by Corollary 1.2, \mathcal{F} embeds, as a K -algebra, in an ultrapower $A^{\mathcal{U}} \simeq A \otimes_K K^{\mathcal{U}}$ (see Corollary 4.3 below). Hence $K^{\mathcal{U}} \mathcal{F}$ is a finite-dimensional $K^{\mathcal{U}}$ -algebra. On the other hand, since \mathcal{F} is residually nilpotent (see, for example, [Ba, §4.2.10] for the case of Lie algebras; the general case is treated identically), the finite-dimensionality of \mathcal{F} would imply nilpotency of \mathcal{F} , and hence nilpotency of A , a contradiction. Consequently, \mathcal{F} is infinite-dimensional over K , and $\mathcal{F} \otimes_K K^{\mathcal{U}}$ is infinite-dimensional over $K^{\mathcal{U}}$. We will use essentially the same type of arguments below in §5, when describing an alternative approach to a result of Kushkulei and Razmyslov about varieties generated by simple finite-dimensional Lie algebras.

3. GROUPS

As the class of (all) groups is, obviously, ideal-determined, Theorem 1.1 and Corollary 1.2 apply to it.

Let us single out an important condition for groups of which the finite subdirect irreducibility is a consequence:

Lemma 3.1. *A group in which any two commuting elements generate a cyclic subgroup either of prime order, or of infinite order, is finitely subdirectly irreducible.*

Proof. Let G be a group with the given property, and N, M be two normal subgroups of G . Take $x \in N, y \in M, x, y \neq 1$. If x, y do not commute, then $1 \neq xyx^{-1}y^{-1} \in N \cap M$. If x, y commute, then they generate a cyclic subgroup of G , generated by a single element $a \in G$, either of prime order p , or of infinite order. Write $x = a^n, y = a^m$ for some $0 < n, m < p$ in the first case, and some nonzero integers n, m in the second one. We have $a^{nm} \in M \cap N$, and due to the restriction on n, m , $a^{nm} \neq 1$ in both cases. \square

Now we can establish Corollary 1.3 for the variety of all groups and for Burnside varieties:

Corollary 3.2 (Criterion for absence of non-trivial identities for groups). *For a group G belonging to one of the following varieties: all groups, groups satisfying the identity $x^p = 1$ for a prime $p \geq 673$, the following are equivalent:*

- (i) G does not satisfy a nontrivial identity within the given variety;
- (ii) any free group in the variety embeds in an ultrapower of G ;
- (iii) any free group in the variety embeds in a group elementarily equivalent to G .

The restriction on p is stipulated, of course, by the celebrated Novikov–Adian solution of the Burnside problem (see [Ad]): the Burnside group $B(n, m)$ of exponent m freely generated by n elements, is infinite for odd $m \geq 665$, and 673 is the next prime after 665.

Proof. Follows from Corollary 1.3, Lemma 3.1, and the fact that the corresponding free groups satisfy the hypothesis of Lemma 3.1: from [Ad, Chapter VI, §3, Theorem 3.3] it follows that under the given restriction on p , any abelian subgroup of the Burnside group $B(n, p)$ is cyclic of order p , and from the Nielsen–Schreier theorem about freeness of subgroups of free groups it follows that any abelian subgroup of an absolutely free group is infinite cyclic. \square

The same remarks as those after Corollary 2.5 apply also in the group case. In particular, condition (ii) is equivalent to the condition that the universal theory of G is contained in the universal theory of any free group. As in the case of Lie and associative algebras, “any free group” in conditions (ii) and (iii) of the corollary, can be replaced by “the free group of rank 2”: an embedding of the free group of countable rank into the free group of rank 2 is well-known in the case of absolutely free groups, and is established in [Š] (see also [At, §3] and references therein for the later alternative approaches) in the case of Burnside groups.

Corollary 3.2 is probably known to experts – at least statements equivalent to the implication (i) \Rightarrow (ii) in the case of the variety of all groups can be found in [Bo, Theorem 1] and [DS, Lemma 6.15]. The proofs there are different and based on the fact that ultraproducts are ω_1 -compact.

Question. *Is there a semigroup property such that the corresponding analogs of the results of this section would hold in the class of all semigroups? The same question for the classes of inverse semigroups and quasigroups.*

Note that neither of these classes is ideal-determined.

4. APPLICATION: PI ALGEBRAS

As an application of this machinery, let us demonstrate how one can handle, in a way different from the traditional approaches, some well-known statements from the theory of associative algebras satisfying polynomial identities (usually called PI algebras).

The Regev celebrated “ $A \otimes B$ ” theorem asserts that the tensor product of two PI algebras A and B is PI (see, for example, [KR, Theorem 5.42]). If we want to prove it using results of §1, we encounter a few difficulties: first, to establish relationship between the ultrapower of the tensor product $(A \otimes B)^{\mathcal{U}}$ and the tensor product of ultrapowers $A^{\mathcal{U}} \otimes B^{\mathcal{U}}$ (perhaps, considering some sort of completed tensor product instead of the usual one may help), and, second, to be able to say something about algebras A and B such that their (possibly completed) tensor product contains a free associative algebra. But at least, in this way we are able to provide an alternative proof of the particular case where one of the tensor factors is finite-dimensional (first established by Procesi and Small in [PS] for the even more particular case where one of the tensor factors is a full matrix algebra; a quantitative refinement (degrees of the corresponding standard identities) of Procesi–Small result was later given in [Do, Theorem 3.2]). This particular case is morally important, as semiprime PI algebras embed in matrix algebras over commutative rings (see, for example, [KR, Remark 1.69]), which essentially reduces the semiprime situation to a finite-dimensional one.

Theorem 4.1 (“Baby Regev’s $A \otimes B$ ”). *The tensor product of two associative algebras, one of them PI and the other finite-dimensional, is PI.*

Lemma 4.2. *Let A, B be algebras defined over a field K , A finite-dimensional. Then, for any ultrafilter \mathcal{U} ,*

$$(A \otimes_K B)^{\mathcal{U}} \simeq A \otimes_K B^{\mathcal{U}}$$

(as K -algebras).

Special cases of this assertion were proved many times in literature – for example, in [NN] for the case where A is an associative full matrix algebra, and in [T, Proposition 25] for the case where both A and B are finite-dimensional, and the proof is standard.

Proof. Let $\{a_1, \dots, a_n\}$ be a basis of A . Obviously, for each K -algebra C , each element of $A \otimes_K C$ can be uniquely represented as $\sum_{k=1}^n a_k \otimes c_k$ for some $c_k \in C$. Define a map

$$\varphi : (A \otimes_K B)^{\mathfrak{J}} \rightarrow A \otimes_K B^{\mathfrak{J}}$$

as follows: for $f \in (A \otimes_K B)^{\mathfrak{J}}$ write $f(i) = \sum_{k=1}^n a_k \otimes b_{ki}$, $i \in \mathfrak{J}$ and define $\varphi(f) \in A \otimes_K B^{\mathfrak{J}}$ as $\sum_{k=1}^n a_k \otimes g_k$, where $g_k \in B^{\mathfrak{J}}$ is defined as $g_k(i) = b_{ki}$, $i \in \mathfrak{J}$. Writing multiplication in A in terms of the basis elements, one can see that φ is an isomorphism of K -algebras.

The ideal $\mathcal{I}((A \otimes_K B)^{\mathfrak{J}}, \mathcal{U})$ maps under φ to $A \otimes_K \mathcal{I}(B^{\mathfrak{J}}, \mathcal{U})$, so factoring out both sides of the isomorphism φ by the corresponding ideals, we get:

$$\begin{aligned} (A \otimes_K B)^{\mathcal{U}} &= (A \otimes_K B)^{\mathfrak{J}} / \mathcal{I}((A \otimes_K B)^{\mathfrak{J}}, \mathcal{U}) \simeq (A \otimes_K B^{\mathfrak{J}}) / (A \otimes_K \mathcal{I}(B^{\mathfrak{J}}, \mathcal{U})) \\ &\simeq A \otimes_K (B^{\mathfrak{J}} / \mathcal{I}(B^{\mathfrak{J}}, \mathcal{U})) = A \otimes_K B^{\mathcal{U}}. \end{aligned}$$

□

Corollary 4.3 ([T, Proposition 21]). *Let A be a finite-dimensional algebra defined over a field K . Then, for any ultrafilter \mathcal{U} , $A^{\mathcal{U}} \simeq A \otimes_K K^{\mathcal{U}}$ (as K -algebras).*

Proof. Put $B = K$.

□

Proof of Theorem 4.1. Let A be a finite-dimensional associative algebra, and B a PI algebra, defined over a field K . Suppose $A \otimes_K B$ is not PI. Then by Corollary 2.5, some ultrapower $(A \otimes_K B)^{\mathcal{U}}$, considered as a $K^{\mathcal{U}}$ -algebra, contains a free associative subalgebra \mathcal{F} of finite rank. By Lemma 4.2,

$$(A \otimes_K B)^{\mathcal{U}} \simeq (A \otimes_K K^{\mathcal{U}}) \otimes_{K^{\mathcal{U}}} B^{\mathcal{U}}$$

as $K^{\mathcal{U}}$ -algebras.

Since \mathcal{F} is finitely-generated, we may choose a finitely-generated $K^{\mathcal{U}}$ -subalgebra B' of $B^{\mathcal{U}}$ such that \mathcal{F} is a subalgebra of $(A \otimes_K K^{\mathcal{U}}) \otimes_{K^{\mathcal{U}}} B'$. Since B is PI, $B^{\mathcal{U}}$ is PI, and B' is PI. The Shirshov height theorem implies that B' has polynomial growth (or, in other words, its Gelfand-Kirillov dimension is finite; see, for example, proof of Theorem 9.19 in [KR]). As $A \otimes_K K^{\mathcal{U}}$ is finite-dimensional (over $K^{\mathcal{U}}$), the tensor product $(A \otimes_K K^{\mathcal{U}}) \otimes_{K^{\mathcal{U}}} B'$ has polynomial growth too. But this contradicts the fact that its subalgebra \mathcal{F} has exponential growth. □

Needless to say, this proof, unlike those in [PS], as well as all the proofs of the full-fledged Regev's $A \otimes B$, is absolutely non-constructive, as it uses existence of an ultrafilter, and, therefore, axiom of choice.

Along the same lines one may treat, at least in some particular cases, a number of other well-known results from PI theory: commutativity of an ordered PI algebra; Piness of a finitely-graded algebra with PI “null component”; of an algebra with a group action whose fixed point subalgebra is PI; of a localization of a PI algebra; of algebras with involution, etc.

Let us note yet another immediate application to a situation which resonates with the universal-algebraic setup of §1. A number of authors considered a property of *definable principal congruences* in algebraic systems. For rings, this amounts to saying that the property that an element x of a ring belongs to the (principal) ideal generated by an element y , is expressed as a first-order formula in free variables x, y of the theory of rings. For an associative ring R , the fact that $\text{Var}(R)$ has definable principal congruences, can be expressed as a certain formula of the universal theory of rings, evidently not satisfied in

free associative rings (see [Si, Lemma 1]). From this and from the ring-theoretic version of Corollary 2.5 immediately follows that if $\text{Var}(R)$ has definable principal congruences, then R is PI, what is the main result of [Si], obtained there with appeal to some deep result from PI theory.

One may try to apply the same reasoning to the known open problem: suppose an associative algebra R is represented as the vector space sum of its subalgebras: $R = A + B$. If A, B are PI, is it true that R is PI? (see [FGL] and [KP] with a transitive closure of references therein). It is easy to see that the operation of taking ultraproduct commutes with the operation of taking the vector space sum: $R^{\mathcal{U}} = A^{\mathcal{U}} + B^{\mathcal{U}}$. Consequently, the question can be reduced to the following one: is it possible that the vector space sum $A + B$ of two PI algebras can contain a free associative subalgebra? If $A + B$ is finitely-generated, the impossibility of this follows from the same growth argument as in the proof of Theorem 4.1. The difficulty, however, lies in the fact that it is not clear how to reduce the situation to a finitely-generated one, as the multiplication between A and B can be intertwined in a complicated way.

5. APPLICATION: ALGEBRAS WITH THE SAME IDENTITIES

In [R, §5] (see also [SZ]), Kushkulei and Razmyslov obtained results stating that some classes of finite-dimensional algebras (e.g., prime over an algebraically closed field) are uniquely determined by their identities. Another result in this direction:

Theorem 5.1. *Let \mathcal{P} be a class of finite-dimensional algebras satisfying the following conditions:*

- (i) *If $A, B \in \mathcal{P}$, A and B are defined over the same field, A is a subalgebra of B , and $\text{Var}(A) = \text{Var}(B)$, then $A = B$.*
- (ii) *\mathcal{P} is closed under elementary equivalence in the first-order two-sorted theory of pairs (algebra over a field, field).*
- (iii) *\mathcal{P} contains all finite-dimensional prime algebras.*

Then \mathcal{P} satisfies the following strengthening of condition (i): if $A, B \in \mathcal{P}$, A and B are defined over the same field, and $\text{Var}(A) = \text{Var}(B)$, then $A \simeq B$.

We stress that the base field over which algebras in the class \mathcal{P} are defined, is not fixed.

Proof. Let $A, B \in \mathcal{P}$, both defined over a field K , be such that $\text{Var}(A) = \text{Var}(B)$. Lemma 2.4, Corollary 1.2 and Corollary 4.3 imply that a free algebra \mathcal{F} in the variety $\text{Var}(A) = \text{Var}(B)$ embeds, as a K -algebra, into an algebra $A \otimes K^{\mathcal{U}}$ for some ultrafilter \mathcal{U} . Hence $K^{\mathcal{U}} \mathcal{F}$ is isomorphic, as a $K^{\mathcal{U}}$ -algebra, to a subalgebra of $A \otimes K^{\mathcal{U}}$. It is easy to see that primeness of the K -algebra \mathcal{F} implies primeness of the $K^{\mathcal{U}}$ -algebra $K^{\mathcal{U}} \mathcal{F}$. By (iii), the latter algebra belong to \mathcal{P} , and by the Łoś theorem and (ii), the $K^{\mathcal{U}}$ -algebra $A \otimes_K K^{\mathcal{U}}$ belongs to \mathcal{P} . Obviously, $K^{\mathcal{U}} \mathcal{F}$ satisfies over $K^{\mathcal{U}}$ the same identities as A over K . Then by (i), $K^{\mathcal{U}} \mathcal{F} \simeq A \otimes_K K^{\mathcal{U}}$. By the same reasoning, $K^{\mathcal{U}} \mathcal{F} \simeq B \otimes_K K^{\mathcal{U}}$. Hence $A \otimes_K K^{\mathcal{U}} \simeq B \otimes_K K^{\mathcal{U}}$ as $K^{\mathcal{U}}$ -algebras, and $A \simeq B$ as K -algebras. \square

This theorem allows, for example, to obtain an alternative proof of results of Kushkulei and Razmyslov in an important particular case of finite-dimensional simple Lie algebras over an algebraically closed field of characteristic zero (see [R, §5, Corollary 1 and Comments]). Indeed, the hypothesis of the theorem is satisfied for such class of algebras: (i) can be proved with the help of the well-known Dynkin's classification [Dy] of semisimple subalgebras of semisimple Lie algebras, (ii) is evident, and (iii) follows from the obvious fact that for finite-dimensional Lie algebras over a field of characteristic zero, simplicity is equivalent to primeness.

The same approach can be applied to finite-dimensional simple Jordan algebras (which follows from the general results of Kushkulei and Razmyslov and also established independently in [DR]), as well as to graded simple associative algebras ([KZ]).

Along the same lines one may treat varieties generated by affine Kac–Moody algebras. Consider, for example, a Lie algebra of the form

$$(7) \quad \widehat{\mathfrak{g}} = (\mathfrak{g} \otimes_K K[t, t^{-1}]) \oplus Kz,$$

where \mathfrak{g} is a split finite-dimensional simple Lie algebra defined over a field K of characteristic zero, $K[t, t^{-1}]$ is the algebra of Laurent polynomials, z is the central element, and the multiplication between elements of $\mathfrak{g} \otimes K[t, t^{-1}]$ is twisted by the well-known 2-cocycle:

$$[x \otimes f, y \otimes g] = [x, y] \otimes fg + (x, y) \text{Res}\left(\frac{df}{dt}g\right)z$$

where $x, y \in \mathfrak{g}$, $f, g \in K[t, t^{-1}]$, and (\cdot, \cdot) is the Killing form on \mathfrak{g} .

In [Za] it is shown, among other, that $\text{Var}(\widehat{\mathfrak{g}}) = \llbracket \text{Var}(\mathfrak{g}), \mathfrak{E} \rrbracket$, where \mathfrak{E} is the variety consisting of the single zero algebra, and $\llbracket \cdot, \cdot \rrbracket$ is the standard commutator of varieties as defined in [Ba, §4.3.8] (in other words, for a variety \mathfrak{V} , $\llbracket \mathfrak{V}, \mathfrak{E} \rrbracket$ is nothing but a variety defined by identities of the form $[f(x_1, \dots, x_n), x_{n+1}] = 0$, where $f(x_1, \dots, x_n) = 0$ is an identity in \mathfrak{V}). Let us complement this result by showing that free algebras in $\text{Var}(\widehat{\mathfrak{g}})$ embed in algebras whose structure closely resembles those of $\widehat{\mathfrak{g}}$.

By [St, Corollary 2.2], every ideal of the Lie algebra $\mathfrak{g} \otimes K[t, t^{-1}]$ is of the form $\mathfrak{g} \otimes I$, where I is an ideal of $K[t, t^{-1}]$. Since $K[t, t^{-1}]$ does not have zero divisors, it is prime, hence $\mathfrak{g} \otimes K[t, t^{-1}]$ is prime, and $\widehat{\mathfrak{g}}$ is a central extension of a prime Lie algebra. By an obvious modification of the proof of Lemma 2.4, one get that for the free algebra \mathcal{L} in $\text{Var}(\widehat{\mathfrak{g}})$ of countable rank, $\mathcal{L}/Z(\mathcal{L})$ is prime, and, by Corollary 1.2 and Lemma 4.2, embeds in $\mathfrak{g} \otimes K[t, t^{-1}]^{\mathcal{U}}$ for some ultrafilter \mathcal{U} . From the results of [Ba, §4.4] it follows that \mathcal{L} embeds in a central extension of $\mathfrak{g} \otimes K[t, t^{-1}]^{\mathcal{U}}$. The latter, by [Ka, Theorem 3.3 and Corollary 3.5], is described in terms of the first-order cyclic homology of $K[t, t^{-1}]^{\mathcal{U}}$, so we get an embedding

$$\mathcal{L} \hookrightarrow \left(\mathfrak{g} \otimes_{K^{\mathcal{U}}} K[t, t^{-1}]^{\mathcal{U}} \right) \oplus HC_1(K[t, t^{-1}]^{\mathcal{U}}).$$

The multiplication in the right-hand side Lie algebra is defined by the formula

$$[x \otimes F, y \otimes G] = [x, y] \otimes FG + (x, y) \overline{F \wedge G},$$

where $x, y \in \mathfrak{g}$, $F, G \in K[t, t^{-1}]^{\mathcal{U}}$, and $\overline{F \wedge G}$ denotes the corresponding homology class in $HC_1(K[t, t^{-1}]^{\mathcal{U}})$.

The addition to (7) of the $Kt \frac{d}{dt}$ term, or twisting by automorphisms of \mathfrak{g} , do not significantly change the picture, and can be treated in the same way.

Similarly, one may treat varieties generated by modular semisimple Lie algebras. According to the classical Block theorem, a typical finite-dimensional semisimple Lie algebra over a field K of characteristic p which is not isomorphic to the sum of simple ones, has the form

$$\left(S \otimes_K K[t_1, \dots, t_n] / (t_1^p, \dots, t_n^p) \right) \oplus \left(1 \otimes_K D \right),$$

where S is a simple Lie algebra, $K[t_1, \dots, t_n] / (t_1^p, \dots, t_n^p)$ is the reduced polynomial algebra, and D is a derivation algebra of $K[t_1, \dots, t_n] / (t_1^p, \dots, t_n^p)$ such that the latter does not have D -invariant ideals.

To such algebras, Lemma 2.4 is applicable, and, as in the Kac–Moody case, further application of Corollary 1.2 and Lemma 4.2 gives an embedding (as $K^{\mathcal{U}}$ -algebras)

$$\mathcal{L} \hookrightarrow \left(S \otimes_{K^{\mathcal{U}}} K^{\mathcal{U}}[t_1, \dots, t_n] / (t_1^p, \dots, t_n^p) \right) \oplus \left(1 \otimes_{K^{\mathcal{U}}} (D \otimes_K K^{\mathcal{U}}) \right)$$

for some ultrafilter \mathcal{U} .

6. APPLICATION: TARSKI’S MONSTERS

Under *Tarski’s monster of type p* , p being a prime (respectively, of type ∞^\dagger) we understand an infinite nonabelian group all whose proper subgroups are cyclic of order p (respectively, of infinite order). Such groups were constructed, among other groups with exotic-looking restrictions on subgroups, by Olshanskii in the framework of his celebrated machinery of geometrically-motivated manipulations with group presentations (see, for example, [O1, Chapter 9, §28.1]). In Olshanskii’s works, the existence of

[†]Added September 16, 2025: The standard name in the literature for what we call “Tarski monster of type ∞ ” is “torsion-free Tarski monster”.

Tarski's monsters of type p is established for $p > 10^{75}$. Later, in [AL] this estimate has been reduced to $p > 1003$.

Let G be a finitely-generated group, and

$$(8) \quad \{1\} \rightarrow \mathcal{N} \rightarrow \mathcal{F} \rightarrow G \rightarrow \{1\}$$

its presentation, where \mathcal{F} is a free group of finite rank, and \mathcal{N} is a normal subgroup of relations. The *girth* of the presentation (8) is the minimal length of elements of \mathcal{N} , i.e., the minimal length of relations between the chosen generators of G (or, in other words, the minimal length of a simple loop in the corresponding Cayley graph). The *girth* of G is the supremum of girths of all its presentations with a finite number of generators. This natural notion was introduced and studied recently by Akhmedov in [Ak1] and [Ak2], and by Schleimer in [Sc]. One of the interesting questions arising in that regard is to construct groups of infinite girth.

As noted in the above-mentioned works, a group satisfying a nontrivial identity cannot have an infinite girth. To circumvent this obstacle, let us introduce the notion of *relative girth* – a girth relative to all identities a group satisfies: in the definition of girth above, replace in (8) the absolutely free group \mathcal{F} by a free group (of finite rank) in the variety $\text{Var}(G)$.

Theorem 6.1.

- (i) A Tarski's monster of type p does not satisfy any nontrivial identity except $x^p = 1$ and its consequences, if and only if it has infinite relative girth.
- (ii) A Tarski's monster of type ∞ does not satisfy any nontrivial identity if and only if it has infinite girth.

Remark (A. Olshanskii). Tarski's monsters satisfying condition (ii) of the theorem do exist (and, moreover, there is an abundance of them), and they can be constructed in the following way.

According to [O2, Corollary 1], each non-cyclic torsion-free hyperbolic group G_0 has a homomorphic image G which is a Tarski's monster of type ∞ . Such monsters are constructed by subsequent applications of [O2, Theorem 2], as the direct limit of a system of surjective maps of groups $G_0 \rightarrow G_1 \rightarrow \dots$. Each G_n is a non-cyclic torsion-free hyperbolic group (and hence contains a free subgroup of countable rank), and is 2-generated for $n \geq 1$. Let us denote, by abuse of notation, the corresponding generators by the same letters a, b (so, $a, b \in G_n$ are images of $a, b \in G_{n-1}$), and each G_n is obtained from G_{n-1} by adding additional relations between these two generators. Also, the injectivity radius of each surjection $G_{n-1} \rightarrow G_n$ (i.e., the maximal number r such that the map is injective on all words of length $\leq r$) can be chosen to be arbitrarily large.

Enumerate all the non-trivial words in the free group of countable rank \mathcal{F} as v_1, v_2, \dots . Since each G_n contains a copy of \mathcal{F} , there are elements in G_n such that the value of v_n on these elements is different from 1. Writing these elements in terms of the generators a, b , we get

$$(9) \quad w_n(a, b) = v_n(w_{n1}(a, b), w_{n2}(a, b), \dots) \neq 1 \text{ in } G_n$$

for some (finite number of) words $w_n, w_{n1}, w_{n2}, \dots$.

Now, on each step choose the injectivity radius of the surjection $G_n \rightarrow G_{n+1}$ larger than the length of all words w_1, \dots, w_n constructed on the previous steps. Consequently, (9) holds in all groups G_{n+1}, G_{n+2}, \dots , and hence in the limit group G . This implies that $v_n = 1$, for any n , cannot be an identity of G .

In [GG], the absence of nontrivial identities in a Tarski's monster G of type ∞ is characterized in terms of an action of the group of outer automorphisms of a free group of rank n on n -tuples of G .

Question. Prove existence of Tarski's monsters satisfying condition (i) of Theorem 6.1.

Proof of Theorem 6.1. The “only if” part is obvious, so let us prove the “if” part. Let G be Tarski's monster either of type p which does not satisfy any nontrivial identity except $x^p = 1$ and its consequences, or of type ∞ which does not satisfy a nontrivial identity. By Corollary 3.2, a group elementarily equivalent to G contains a subgroup isomorphic to a relatively free subgroup \mathcal{G} of rank 2 (which is the free Burnside

group $B(2, p)$ in the case (i), or the free group in the case (ii)). Let x, y be the free generators of \mathcal{G} , and

$$\{w_1(x, y) = x, w_2(x, y) = x^{-1}, w_3(x, y) = y, w_4(x, y) = y^{-1}, \dots, w_{k_n}(x, y)\}$$

the set of all words of \mathcal{G} of length $\leq n$. The existence of this “initial piece of \mathcal{G} of length n ” can be written as the first-order property:

$$\exists x \exists y : \bigwedge_{1 \leq i < j \leq k_n} w_i(x, y) \neq w_j(x, y).$$

Consequently, for each $n \in \mathbb{N}$, the same first-order formula holds in G ; let $x_n, y_n \in G$ be the corresponding elements. Obviously, x_n, y_n do not commute except, possibly, for some small values of n , and, therefore, generate G . This provides a presentation of G of (relative) girth $> n$. \square

Note another interesting consequence of Theorem 6.1.

The *growth sequence* of a group G is a sequence whose n th term equal to the (minimal) number of generators of the n th fold direct power of G . See [W] for a brief history of the subject and further references. In particular, in a number of works, including [W], a considerable effort was put into construction of groups whose growth sequence is constant, each term is equal to 2. Theorem 6.1 provides further such examples, in view of the following general elementary fact:

Lemma 6.2. *If a finitely-generated simple group G has infinite relative girth, then its growth sequence is constant, each term is equal to the minimal number of generators of G .*

Proof. Let n be the minimal number of generators of G . Infinity of the relative girth of G means that there is an infinite sequence $\mathcal{N}_1, \mathcal{N}_2, \dots$ of normal subgroups of the free group \mathcal{G} in $\text{Var}(G)$ of rank n such that for every i the length of each word in \mathcal{N}_i is $\geq i$, and $\mathcal{G}/\mathcal{N}_i \simeq G$.

Let us prove by induction that for each $k \in \mathbb{N}$ there is a sequence $i_1 < i_2 < \dots < i_k$ such that

$$(10) \quad \mathcal{G}/(\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k}) \simeq G \times G \times \dots \times G \quad (k \text{ times}).$$

For $k = 1$ we may take $i_1 = 1$. Suppose that for some $k > 1$ the isomorphism (10) holds. It is obvious that $\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k} \neq \{1\}$. On the other hand, since $\bigcap_{i > i_k} \mathcal{N}_i = \{1\}$, there is $i_{k+1} > i_k$ such that

$$\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k} \not\subseteq \mathcal{N}_{i_{k+1}}.$$

Since G is simple, $\mathcal{N}_{i_{k+1}}$ is a maximal normal subgroup in \mathcal{G} , and

$$\mathcal{G} = (\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k}) \mathcal{N}_{i_{k+1}}.$$

Then:

$$\begin{aligned} \mathcal{G}/(\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k} \cap \mathcal{N}_{i_{k+1}}) &\simeq \mathcal{G}/(\mathcal{N}_{i_1} \cap \dots \cap \mathcal{N}_{i_k}) \times \mathcal{G}/\mathcal{N}_{i_{k+1}} \\ &\simeq G \times \dots \times G \quad (k+1 \text{ times}). \end{aligned}$$

It follows from (10) that each finite direct power of G is n -generated. \square

In fact, nothing in this proof is specific to groups: the corresponding statement can be formulated for general algebraic systems; in particular, it holds also for algebras.

Lemma 6.2 implies that the growth sequence of Tarski’s monsters satisfying conditions of Theorem 6.1 is constant, each term is equal to 2. Note that in [GG, Theorem 8] it is proved, by different methods, that each term in the growth sequence of *any* Tarski’s monster is ≤ 3 , and for any Tarski’s monster of type ∞ it is equal to 2.

7. HOMOMORPHISMS OF DIRECT PRODUCTS

The following is, in a sense, dual version of Theorem 1.1, with embeddings replaced by surjective homomorphisms, with, essentially, the same proof[†].

[†]Added August 22, 2015: the proof of Theorem 7.1 is in error. For a corrected statement, with a different proof, see a sequel to this article, *On the utility of Robinson–Amitsur ultrafilters. II*.

Theorem 7.1. *Let $\{B_i\}_{i \in \mathcal{I}}$ be a set of algebraic systems from an ideal-determined class. If a finitely subdirectly irreducible algebraic system A is a surjective homomorphic image of the direct product $\prod_{i \in \mathcal{I}} B_i$, then there is an ultrafilter \mathcal{U} on the set \mathcal{I} such that A is a surjective homomorphic image of the ultraproduct $\prod_{\mathcal{U}} B_i$.*

Proof. Let $\alpha : \prod_{i \in \mathcal{I}} B_i \rightarrow A$ be a given surjective homomorphism. Define

$$\mathcal{S} = \{\{i \in \mathcal{I} \mid f(i) \neq e\} \mid f \in \prod_{i \in \mathcal{I}} B_i, \alpha(f) \neq e\}.$$

Let us verify that intersection of any two elements of \mathcal{S} contains an element of \mathcal{S} . Let $S, T \in \mathcal{S}$, say, $S = \{i \in \mathcal{I} \mid f(i) \neq e\}$ and $T = \{i \in \mathcal{I} \mid g(i) \neq e\}$ for some $f, g \in \prod_{i \in \mathcal{I}} B_i$ not vanishing under α . Since A is finitely subdirectly irreducible, it contains an element $a \neq e$ belonging to the intersection of ideals generated by $\{\alpha(f)\}$ and $\{\alpha(g)\}$. Take some element $u \in \prod_{i \in \mathcal{I}} B_i$ from the preimage $\alpha^{-1}(a)$. Let $i \in \mathcal{I}$ such that $f(i) = e$. Since $\alpha(u) = a = t(b_1, \dots, b_n, \alpha(f), \dots, \alpha(g))$ for some ideal term t and $b_1, \dots, b_n \in A$, we have $u = t(h_1, \dots, h_n, f, \dots, f)$ for some $h_1 \in \alpha^{-1}(b_1), \dots, h_n \in \alpha^{-1}(b_n)$, and

$$\begin{aligned} u(i) &= t(h_1, \dots, h_n, f, \dots, f)(i) = t(h_1(i), \dots, h_n(i), f(i), \dots, f(i)) \\ &= t(h_1(i), \dots, h_n(i), e, \dots, e) = e. \end{aligned}$$

Coupling this with a similar assertion for g , we get that

$$S \cap T \supset \{i \in \mathcal{I} \mid u(i) \neq e\} \in \mathcal{S}.$$

Thus \mathcal{S} satisfies the finite intersection property and is contained in some ultrafilter \mathcal{U} on \mathcal{I} . If $f \in \mathcal{I}\left(\prod_{i \in \mathcal{I}} B_i, \mathcal{U}\right)$, then $\{i \in \mathcal{I} \mid f(i) = e\} \in \mathcal{U}$, and, since \mathcal{U} is ultrafilter, $\{i \in \mathcal{I} \mid f(i) \neq e\} \notin \mathcal{U}$, and hence $\{i \in \mathcal{I} \mid f(i) \neq e\} \notin \mathcal{S}$ and $\alpha(f) = e$. This shows that $\mathcal{I}\left(\prod_{i \in \mathcal{I}} B_i, \mathcal{U}\right)$ lies in $\text{Ker } \alpha$, and the map α factors through the ultraproduct $\left(\prod_{i \in \mathcal{I}} B_i\right) / \mathcal{I}\left(\prod_{i \in \mathcal{I}} B_i, \mathcal{U}\right) = \prod_{\mathcal{U}} B_i$. \square

In [Zi, §5.3], it is asked whether an ultraproduct of finite groups can be mapped surjectively to a compact simple Lie group G , in particular, $\text{SO}(3)$. According to Theorem 7.1, this is equivalent to the question whether a direct product of finite groups can be mapped surjectively to G .

Question. *What can be said about the class of groups obtained by closure of finite groups with respect to direct products and homomorphic images? Which simple groups, matrix groups, Lie groups, lie in this class?*

A question when a homomorphic image of a direct product of groups or algebras can be factored through an ultraproduct, or, more generally, through a product of a finite number of ultraproducts, was studied also in the recent interesting papers [Be] and [BN]. The emphasis there is, however, different: one imposes various conditions (such as simplicity, solvability, or nilpotency) not on A but on B_i 's.

8. FURTHER SPECULATIONS

Here we indicate some of our initial reasons for looking into all this, of a highly speculative character.

8.1. Lie-algebraic monsters. A problem of existence of Lie-algebraic analogs of Tarski's monsters – namely, of infinite-dimensional Lie algebras all whose proper subalgebras are one-dimensional – is, arguably, one of the most difficult problems in the abstract theory of infinite-dimensional Lie algebras (see, for example, [G] and [L]). For such hypothetical Lie algebras which do not satisfy a nontrivial identity, an analog of Theorem 6.1 would hold.

Question. *Study the notion of (relative) girth for Lie algebras.*

A simpler, but still open problem concerns *finite-dimensional* Lie algebras all whose proper subalgebras are one-dimensional (called L_1 -algebras in the sequel). Over perfect fields, every L_1 -algebra has dimension 2 or 3. In characteristic zero this is an immediate consequence of simple reasonings about

forms of algebras and the classical structure theory, and in the case of the perfect base field of characteristic $p > 0$ this follows from the powerful results of Premet [Pr1] (for $p > 5$) and [Pr2] (for $p = 5$). See, for example, [G, Proposition 1] or [L, Proposition 3.1] for details.

Over non-perfect fields, we do not have the usual tools of Galois theory to handle forms of algebras, as in [Pr1], and, to quote [V], “the existence [of an L_1 -algebra] of dimension greater than 3 is an interesting open problem”. Using reasonings similar to those in §2, it is possible to prove that a prime finite-dimensional Lie algebra over an infinite field, all whose abelian subalgebras are 1-dimensional, satisfies a functional identity of the form

$$(11) \quad \lambda(x, y)u(x, y) + \mu(x, y)v(x, y) = 0,$$

where u and v are certain Lie words in 2 variables, and λ and μ are elements of the base field, depending, in general, on the elements x and y of the algebra. We conjecture that no simple finite-dimensional Lie algebra of dimension > 3 can satisfy a nontrivial functional identity of the form (11). If this conjecture is true, it will prove the absence of L_1 -algebras of dimension > 3 over an arbitrary base field.

8.2. The Tits alternative. Another reason was the desire to provide an alternative proof of the celebrated Tits alternative, or for one of its not less celebrated consequences, such as growth dichotomy for linear group. The Tits alternative claims that a linear group contains either a solvable subgroup of finite index, or a nonabelian free subgroup. A nice and important, yet admitting a few-lines elementary proof, result of Platonov [Pl] states that a linear group which has a nontrivial identity, contains a solvable subgroup of finite index. Modulo this result, the proof of the Tits alternative reduces to establishing that a linear group G which does not satisfy a nontrivial identity, contains a nonabelian free subgroup. One may naively argue as follows: by Corollary 3.2, a group, elementarily equivalent to G , contains a nonabelian free subgroup. From this we may infer some first-order properties of G , for example, that it contains “a piece of a nonabelian free group of arbitrarily large length”, like in the proof of Theorem 6.1. On the other hand, as linearity is a sort of finiteness condition, one may hope that these first-order properties may help to construct a nonabelian free subgroup in G . If successful, this approach would provide a proof of the Tits alternative drastically different from all the proofs given so far.

8.3. Jacobson’s problem. An old open problem due to Jacobson asks whether a Lie p -algebra L such that for every $x \in L$ there is $n(x) \in \mathbb{N}$ satisfying

$$(12) \quad x^{p^{n(x)}} = x,$$

is abelian? As one of the first steps, one may wish to prove that such Lie algebras satisfy a nontrivial identity; or, for example, that there are no simple (or even prime) such algebras. In both of these cases, by Lemma 2.4 and Corollary 1.2, a free Lie algebra embeds in an algebra elementarily equivalent to L , over some elementary extension of the base field. The condition (12) is not the first-order property (unless all $n(x)$ are bounded, in which case the problem is trivial), but one may hope to derive from it some first-order consequences which will come into contradiction with the existence of a free Lie subalgebra.

ACKNOWLEDGEMENTS

Thanks are due to Shelly Garion, Misha Gavrillovich, Sergei Ivanov, Ruvim Lipyanski, Alexander Olshanskii, Mark Sapir, Ivan Shestakov, Irina Sviridova, Jenő Szegedi, and Boris Zilber for helpful and interesting comments.

A previous version of this manuscript was rejected by *Journal of the London Mathematical Society*, but I owe to the anonymous referee of that submission a great deal of improvements and corrections, both explicit and implicit: in particular, he gave me an idea to use the condition of finite subdirect irreducibility, and to formulate Theorem 1.1 in the language of universal algebra. The final touches were done during my visit to IHÉS at February 2012.

This work was partially supported by grants ERMOS7 (Estonian Science Foundation and Marie Curie Actions) and ETF9038 (Estonian Science Foundation).

REFERENCES

- [Ad] S.I. Adian, *The Burnside Problem and Identities in Groups*, Nauka, Moscow, 1975 (in Russian); Springer, 1979 (English translation).
- [AL] ——— and I.G. Lysenok, *On groups all of whose proper subgroups are finite cyclic*, *Izv. Akad. Nauk SSSR Ser. Mat.* **55** (1991), 933–990 (in Russian); *Math. USSR Izv.* **39** (1992), 905–957 (English translation).
- [Ak1] A. Akhmedov, *On the girth of finitely generated groups*, *J. Algebra* **268** (2003), 198–208.
- [Ak2] ———, *The girth of groups satisfying Tits alternative*, *J. Algebra* **287** (2005), 275–282.
- [Am1] S.A. Amitsur, *Generalized polynomial identities and pivotal monomials*, *Trans. Amer. Math. Soc.* **114** (1965), 210–226; reprinted in *Selected Papers of S.A. Amitsur with Commentary*, Part 1, AMS, 2001, 191–208.
- [Am2] ———, *Prime rings having polynomial identities with arbitrary coefficients*, *Proc. London Math. Soc.* **17** (1967), 470–486; reprinted in *Selected Papers of S.A. Amitsur with Commentary*, Part 1, AMS, 2001, 417–433.
- [At] V.S. Atabekyan, *On normal subgroups in the periodic products of S.I. Adian*, *Trudy Mat. Inst. Steklova* **274** (2011), 15–31 (in Russian); *Proc. Steklov Inst. Math.* **274** (2011), 9–24 (English translation).
- [Ba] Yu.A. Bahturin, *Identical Relations in Lie Algebras*, Nauka, Moscow, 1985 (in Russian); VNU Sci. Press, Utrecht, 1987 (English translation).
- [BS] J.L. Bell and A.B. Slomson, *Models and Ultraproducts*, 2nd revised printing, North-Holland, 1971.
- [Be] G.M. Bergman, *Families of ultrafilters, and homomorphisms on infinite direct product algebras*; arXiv:1301.6383v1.
- [BN] ——— and N. Nahlus, *Homomorphisms on infinite direct product algebras, especially Lie algebras*, *J. Algebra* **333** (2011), 67–104; arXiv:0910.5181.
- [Bo] M. Boffa, *Elimination of inverses in groups*, *Model Theory of Groups and Automorphism Groups* (ed. D.M. Evans), London Math. Soc. Lect. Note Ser. **244** (1997), 134–143.
- [CEL] I. Chajda, G. Eigenthaler, and H. Länger, *Congruence Classes in Universal Algebra*, Heldermann Verlag, 2003.
- [C] P.M. Cohn, *Universal Algebra*, Harper & Row, 1965.
- [Do] M. Domokos, *On algebras satisfying symmetric identities*, *Arch. Math.* **63** (1994), 407–413.
- [DR] V.S. Drensky and M.L. Racine, *Distinguishing simple Jordan algebras by means of polynomial identities*, *Comm. Algebra* **20** (1992), 309–327.
- [DS] C. Druţu and M. Sapir (with an appendix by D. Osin and M. Sapir), *Tree-graded spaces and asymptotic cones of groups*, *Topology* **44** (2005), 959–1058; arXiv:math/0405030.
- [Dy] E.B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, *Mat. Sbornik* **30** (1952), 349–462 (in Russian); *AMS Transl.* **6** (1957), 111–244 (English translation); reprinted in *Selected Papers of E.B. Dynkin with Commentary*, AMS, 2000, 175–244.
- [E] P. Eklof, *Ultraproducts for algebraists*, *Handbook of Mathematical Logic* (ed. J. Barwise), North-Holland, 1977, 105–137.
- [FGL] B. Felzenszwalb, A. Giambruno, and G. Leal, *On rings which are sums of two PI-subrings: a combinatorial approach*, *Pacific J. Math.* **209** (2003), 17–30.
- [FM] R. Freese and R. McKenzie, *Commutator Theory for Congruence Modular Varieties*, London Math. Soc. Lect. Note Ser. **125** (1987).
- [GG] S. Garion and Y. Glasner, *Highly transitive actions of $\text{Out}(F_n)$* , *Groups, Geometry, and Dynamics* **7** (2013), 357–376; arXiv:1008.0563.
- [G] A.G. Gein, *The modular law and relative complements in the lattice of subalgebras of a Lie algebra*, *Izv. VUZ Mat.* 1987, *N*3, 18–25 (in Russian); *Soviet Math. (Izv. VUZ)* **31** (1987), *N*3, 22–32 (English translation).
- [KR] A. Kanel-Belov and L.H. Rowen, *Computational Aspects of Polynomial Identities*, AK Peters, 2005.
- [Ka] C. Kassel, *Kähler differentials and coverings of complex semisimple Lie algebras extended over a commutative algebra*, *J. Pure Appl. Algebra* **34** (1984), 265–275.
- [KP] M. Kępczyk and E.R. Puczyłowski, *On the structure of rings which are sums of two subrings*, *Arch. Math.* **83** (2004), 429–436.
- [KZ] P. Koshlukov and M. Zaicev, *Identities and isomorphisms of graded simple algebras*, *Lin. Algebra Appl.* **432** (2010), 3141–3148.
- [Ku] A.G. Kurosh, *Lectures on General Algebra*, 2nd ed., Nauka, Moscow, 1973 (in Russian); Pergamon Press, 1965 (English translation of the 1st ed.).
- [L] A.A. Lashkhi, *Lattices with modular identity and Lie algebras*, *Itogi Nauki i Tekhniki. Sovremennye Problemy Matematiki. Noveishie Dostizheniya* **26** (1985), 213–257 (in Russian); *J. Soviet Math.* **38** (1987), 1823–1853 (English translation).
- [MM] L. Makar-Limanov and P. Malcolmson, *Free subalgebras of enveloping fields*, *Proc. Amer. Math. Soc.* **111** (1991), 315–322.
- [M] A.I. Malcev, *Algebraic Systems*, Izdat. Fiz.-Mat. Lit., Moscow, 1970 (in Russian); Springer, 1973 (English translation).
- [NN] A. Niţă and C. Niţă, *Sur l’ultraproduit des anneaux de matrices*, *Bull. Math. Soc. Sci. Math. Rep. Soc. Roumanie* **29** (1985), 275–278.

- [O1] A.Yu. Olshanskii, *The Geometry of Defining Relations in Groups*, Nauka, Moscow, 1989 (in Russian); Kluwer, 1991 (English translation).
- [O2] ———, *On residualizing homomorphisms and G -subgroups of hyperbolic groups*, Intern. J. Algebra Comput. **3** (1993), 365–409.
- [Pl] V.P. Platonov, *Linear groups with identity relations*, Doklady Akad. Nauk BSSR **11** (1967), 581–582 (in Russian).
- [Pr1] A.A. Premet, *Lie algebras without strong degeneration*, Mat. Sbornik **129** (1986), 140–153 (in Russian); Math. USSR Sbornik **57** (1987), 151–163 (English translation).
- [Pr2] ———, *Inner ideals in modular Lie algebras*, Vesti Akad. Navuk BSSR Ser. Fiz.-Mat. Navuk 1986, *N*5, 11–15 (in Russian).
- [PS] C. Procesi and L. Small, *Endomorphism rings of modules over PI-algebras*, Math. Z. **106** (1968), 178–180.
- [R] Yu.P. Razmyslov, *Identities of Algebras and Their Representations*, Nauka, Moscow, 1989 (in Russian); AMS, 1994 (English translation).
- [Sc] S. Schleimer, *On the girth of groups*, Manuscript, 2000;
<http://math.rutgers.edu/~saulsch/Maths/girth.pdf>
- [SZ] I. Shestakov and M. Zaicev, *Polynomial identities of finite dimensional simple algebras*, Comm. Algebra **39** (2011), 929–932.
- [Si] G.E. Simons, *Varieties of rings with definable principal congruences*, Proc. Amer. Math. Soc. **87** (1983), 397–402.
- [Š] V.L. Širvanjan, *Embedding the group $B(\infty, n)$ in the group $B(2, n)$* , Izv. Akad. Nauk SSSR Ser. Mat. **40** (1976), 190–208 (in Russian); Math. USSR Izv. **124** (1976), 161–177 (English translation).
- [St] I. Stewart, *Tensorial extensions of central simple algebras*, J. Algebra **25** (1973), 1–14.
- [T] F. Taha, *Algèbres simples centrales sur les corps ultra-produits de corps p -adiques*, Séminaire d’Algèbre Paul Dubreil et Marie-Paul Malliavin (Paris 1981), Lect. Notes Math. **924** (1982), 89–128.
- [V] V.R. Varea, *Supersimple and upper semimodular Lie algebras*, Comm. Algebra **23** (1995), 2323–2330.
- [W] D.T. Wise, *The rank of a direct power of a small-cancellation group*, Geom. Dedicata **94** (2002), 215–223.
- [Za] M.V. Zaicev, *Varieties and identities of affine Kac–Moody algebras*, Methods in Ring Theory (ed. V.S. Drensky, A. Giambruno, and S. Sehgal), Marcel Dekker, Lect. Notes Pure Appl. Math. **198** (1998), 303–314.
- [Zi] B. Zilber, *Perfect infinities and finite approximation*, Manuscript, November 2012;
<http://people.maths.ox.ac.uk/zilber/inf-to-finite.pdf>

Email address: pasha.zusmanovich@gmail.com