

On extensions of ternary maps to superposition of binary maps

Pasha Zusmanovich

University of Ostrava

June 24, 2015

(based on arXiv:1408.2982)

The last question of Banach

Andrzej Alexiewicz's diary, December 29, 1944:

"There exists a nontrivial example of ternary multiplication, which is not generated by binary multiplication ([Banach](#)). Can any finite set with ternary commutative multiplication be extended so that ternary multiplication is generated by a binary multiplication?"



What does this question mean?

Interpretation of “multiplication” and “generated”?

Connection with:

- ▶ History
- ▶ Semigroups
- ▶ Theory of clones
- ▶ Hilbert's 13th problem (superposition of functions)
- ▶ Logic
- ▶ Combinatorics
- ▶ Computer calculations
- ▶ Lie theory

First interpretation: semigroups

“Multiplication”: ternary semigroup, i.e.

$$f : X \times X \times X \rightarrow X$$

$$f(f(x, y, z), u, v) = f(x, f(y, z, u), v) = f(x, y, f(z, u, v)).$$

Commutativity:

$$f(x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}) = f(x_1, x_2, x_3)$$

for any $\sigma \in S_3$.

“Generation”: $f(x, y, z) = (x * y) * z$.

Theorem (Łoś 1955, Monk–Sioson 1966)

Every (commutative) ternary semigroup can be extended to a (commutative) ternary semigroup given by

$$f(x, y, z) = (x * y) * z,$$

where $*$ is a (commutative) binary semigroup.

Second interpretation: clones

“Multiplication”: an arbitrary map $X \times X \times X \rightarrow X$.

“Generation”: superposition.

Theorem (Sierpiński 1935, Banach 1935)

Any countable number of unary maps on an infinite set can be generated by two maps.

(\Leftrightarrow A countable transformation semigroup is 2-generated).

Generalizations:

- ▶ Other kinds of semigroups, generation “up to approximation” (Schreier–Ulam, Jarnik–Knichal, et al.)
- ▶ Unary \rightarrow multiary

Second interpretation: clones

Theorem

Any countable number of maps of arbitrary arity on a set X can be generated by one binary map on X .

(\Leftrightarrow The clone of all maps on X is generated by its binary fragment).

Webb 1935: finite X (generalization of Sheffer's stroke)

$$p \uparrow q = \neg(p \wedge q)$$

Łoś 1950, Goldstern 2012: infinite X

Conclusion

“Banach's claim” is false in this context.

Interlude

Erdős (on another occasion, from a preface to the *Scottish Book*):

“Now it frequently happens in problems of this sort that the infinite dimensional case is easier to settle than the finite dimensional analogues. This moved **Ulam** and me to paraphrase a well known maxim of the American armed forces in WWII: ‘The difficult we do immediately, the impossible takes a little longer’, viz: ‘The infinite we do immediately, the finite takes a little longer’.”



Stanisław Ulam
(PhD Lvov 1933)



Jerzy Łoś
(student at Lvov, 1937–1939)



Wacław Sierpiński
(professor at Lvov, 1908–1914)

Connection: Hilbert's 13th problem

Theorem (Pólya and Szegő, *Problems and Theorems in Analysis*, 1925, Problem 119 “Are there actually functions of 3 variables?”, Sierpiński 1934,1945)

For any binary bijection $g : X \times X \rightarrow X$ on an (infinite) set X , and any ternary map $f : X \times X \times X \rightarrow X$, there is a binary map $h : X \times X \rightarrow X$ such that

$$f(x, y, z) = g(h(x, y), z).$$

Theorem (Kolmogorov 1956, Arnold 1957)

Any continuous real function in any number of variables can be represented as superposition of continuous real functions in 2 variables.

Questions (Mark Kac–Ulam 1960,1968)

Extensions of Hilbert's 13th problem: on \mathbb{R}^n , in various classes (smooth, analytic).

Connection: Logic

Łukasiewicz's 3-valued logic: "implication" \mathcal{L} :

	0	$\frac{1}{2}$	1
0	1	1	1
$\frac{1}{2}$	$\frac{1}{2}$	1	1
1	0	$\frac{1}{2}$	1

and "negation" N :

$$0 \mapsto 1, \frac{1}{2} \mapsto \frac{1}{2}, 1 \mapsto 0.$$

Theorem (Słupecki 1936)

There are multiary maps on $\{0, \frac{1}{2}, 1\}$ which are not superposition of \mathcal{L} and N .

$2\frac{1}{2}$ th interpretation: clones with additional structure

“Multiplication”: a map $X \times X \times X \rightarrow X$ preserving an additional structure on X .

“Generation”: composition.

“Banach’s claim” is true for smooth functions. For example:

$$F(x, y, z) = f(g(h(x, y), z), z)$$

satisfies the differential equation

$$\frac{\partial^2 F}{\partial x \partial z} \frac{\partial F}{\partial y} - \frac{\partial F}{\partial x} \frac{\partial^2 F}{\partial y \partial z} = 0.$$

Not suitable for finite X .

Third interpretation: operadic-like

“Multiplication”: an arbitrary map $f : X \times X \times X \rightarrow X$.

“Generation”: $f(x, y, z) = (x * y) * z$ or $x * (y * z)$.

For X finite, “Banach’s claim” is true:

number of ternary maps generated by binary ones $< 2|X|^{|X|^2}$

number of ternary maps $= |X|^{|X|^3}$.

Question: number of ternary maps generated by binary ones

$ X = n$	number of binary maps $= n^{n^2}$	number of maps $(x * y) * z$	number of maps $(x * y) * z$ and $x * (y * z)$	number of commutative maps $(x * y) * z$
1	1	1	1	1
2	16	14	21	5
3	19683	19292	38472	48

Absent in OEIS!

Jacobson answers Banach

Theorem

Any ternary map (on a finite set) can be extended to a ternary map generated by a binary map (on a finite set) .

Proof (Jacobson, *Lie and Jordan triple systems*, 1949)

1. Adjoin a “neutral” element e :

$$f(e, X, X) = f(X, e, X) = f(X, X, e) = e.$$

2. $Y = X \times \{m_{x,y} : X \rightarrow X; z \mapsto f(x, y, z)\}$.

3. Binary multiplication on Y :

$$(x, g) * (y, h) = (g(y), m_{x,y}).$$

4. (X, f) is embedded into $(Y, *)$ via $x \mapsto (x, m_{e,e})$.

Modification of this: commutative version, result of Łoś and Monk-Sioson, generalization to n -ary maps, etc., etc.

To je vše. Děkuji.

Slides at <http://justpasha.org/math/banach-ostrava.pdf>